

New England Biolabs Certificate of Analysis


Product Name: NEB® 5-alpha Competent *E. coli* (High Efficiency)
Catalog Number: C2987H
Lot Number: 10051920
Expiration Date: 08/2020
Storage Temperature: -80°C
Specification Version: PS-C2987H/I v1.0

NEB® 5-alpha Competent <i>E. coli</i> (High Efficiency) Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
N3041AVIAL	pUC19 Vector	10047676	Pass
C2987HVIAL	NEB® 5-alpha Competent <i>E. coli</i> (High Efficiency)	10046254	Pass
B9020SVIAL	SOC Outgrowth Medium	10045021	Pass


Assay Name/Specification	Lot # 10051920
Antibiotic Sensitivity (Streptomycin) 15 µl of untransformed NEB® 5-alpha Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate containing Streptomycin will not form colonies after incubation for 16 hours at 37°C.	Pass
Antibiotic Sensitivity (Tetracycline) 15 µl of untransformed NEB® 5-alpha Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate containing Tetracycline will not form colonies after incubation for 16 hours at 37°C.	Pass
Blue-White Screening (α-complementation, Competent Cells) NEB® 5-alpha Competent <i>E. coli</i> (High Efficiency) were shown to be suitable for blue/white screening by α-complementation of the β-galactosidase gene using pUC19.	Pass
Phage Resistance (φ 80) 15 µl of untransformed NEB® 5-alpha Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate does not support plaque formation by phage φ 80 after incubation for 16 hours at 37°C.	Pass
Transformation Efficiency 50 µl of NEB® 5-alpha Competent <i>E. coli</i> (High Efficiency) cells were transformed with 100 pg of pUC19 DNA using the transformation protocol provided. Incubation overnight on LB-Ampicillin plates at 37°C resulted in >1 x 10 ⁹ cfu/µg of DNA.	Pass

Assay Name/Specification	Lot # 10051920
<p>Antibiotic Sensitivity (Ampicillin) 15 µl of untransformed NEB® 5-alpha Competent E. coli (High Efficiency) streaked onto a Rich Broth plate containing Ampicillin will not form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p>Antibiotic Sensitivity (Chloramphenicol) 15 µl of untransformed NEB® 5-alpha Competent E. coli (High Efficiency) streaked onto a Rich Broth plate containing Chloramphenicol will not form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p>Antibiotic Sensitivity (Kanamycin) 15 µl of untransformed NEB® 5-alpha Competent E. coli (High Efficiency) streaked onto a Rich Broth plate containing Kanamycin will not form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p>Antibiotic Sensitivity (Nitrofurantoin) 15 µl of untransformed NEB® 5-alpha Competent E. coli (High Efficiency) streaked onto a Rich Broth plate containing Nitrofurantoin will not form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p>Antibiotic Sensitivity (Spectinomycin) 15 µl of untransformed NEB® 5-alpha Competent E. coli (High Efficiency) streaked onto a Rich Broth plate containing Spectinomycin will not form colonies after incubation for 16 hours at 37°C.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.



Lixin An
Production Scientist
28 Jun 2019



Nick Privitera
Packaging Quality Control Inspector
02 Aug 2019