

New England Biolabs Certificate of Analysis

Product Name: NEB® 5-alpha Competent *E. coli* (High Efficiency)
 Catalog Number: C2987H
 Packaging Lot Number: 10180226
 Expiration Date: 02/2024
 Storage Temperature: -80°C
 Specification Version: PS-C2987H/I v1.0

NEB® 5-alpha Competent <i>E. coli</i> (High Efficiency) Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
N3041AVIAL	pUC19 Vector	10173306	Pass
C2987HVIAL	NEB® 5-alpha Competent <i>E. coli</i> (High Efficiency)	10167802	Pass
B9020SVIAL	SOC Outgrowth Medium	10166777	Pass

Assay Name/Specification	Lot # 10180226
<p>Antibiotic Sensitivity (Kanamycin) 15 µl of untransformed NEB® 5-alpha Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate containing Kanamycin will not form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p>Antibiotic Sensitivity (Nitrofurantoin) 15 µl of untransformed NEB® 5-alpha Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate containing Nitrofurantoin will not form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p>Antibiotic Sensitivity (Streptomycin) 15 µl of untransformed NEB® 5-alpha Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate containing Streptomycin will not form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p>Antibiotic Sensitivity (Tetracycline) 15 µl of untransformed NEB® 5-alpha Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate containing Tetracycline will not form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p>Blue-White Screening (α-complementation, Competent Cells) NEB® 5-alpha Competent <i>E. coli</i> (High Efficiency) were shown to be suitable for blue/white screening by α-complementation of the β-galactosidase gene using pUC19.</p>	Pass


Assay Name/Specification	Lot # 10180226
<p>Phage Resistance (ϕ 80) 15 μl of untransformed NEB® 5-alpha Competent E. coli (High Efficiency) streaked onto a Rich Broth plate does not support plaque formation by phage ϕ 80 after incubation for 16 hours at 37°C.</p>	Pass
<p>Antibiotic Sensitivity (Spectinomycin) 15 μl of untransformed NEB® 5-alpha Competent E. coli (High Efficiency) streaked onto a Rich Broth plate containing Spectinomycin will not form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p>Antibiotic Sensitivity (Chloramphenicol) 15 μl of untransformed NEB® 5-alpha Competent E. coli (High Efficiency) streaked onto a Rich Broth plate containing Chloramphenicol will not form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p>Antibiotic Sensitivity (Chloramphenicol) 15 μl of untransformed NEB® 5-alpha Competent E. coli (High Efficiency) streaked onto a Rich Broth plate containing Chloramphenicol will not form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p>Antibiotic Sensitivity (Ampicillin) 15 μl of untransformed NEB® 5-alpha Competent E. coli (High Efficiency) streaked onto a Rich Broth plate containing Ampicillin will not form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p>Transformation Efficiency 50 μl of NEB® 5-alpha Competent E. coli (High Efficiency) cells were transformed with 100 pg of pUC19 DNA using the transformation protocol provided. Incubation overnight on LB-Ampicillin plates at 37°C resulted in $>1 \times 10^9$ cfu/μg of DNA.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Chris Blanchette
Production Scientist
04 Nov 2022



Nick Privitera
Packaging Quality Control Inspector
10 Feb 2023