

New England Biolabs Certificate of Analysis

Product Name: 5' DNA Adenylation Kit
 Catalog Number: E2610L
 Concentration: 50 μ M
 Packaging Lot Number: 10053873
 Expiration Date: 07/2021
 Storage Temperature: -20°C
 Specification Version: PS-E2610S/L v1.0

5' DNA Adenylation Kit Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
N0757AVIAL	Adenosine 5' Triphosphate	10049239	Pass
M2611AAVIAL	Mth RNA Ligase	10049237	Pass
B2610SVIAL	5' DNA Adenylation Reaction Buffer	10042923	Pass

Assay Name/Specification	Lot # 10053873
<p>Protein Purity Assay (SDS-PAGE) Mth RNA Ligase is \geq 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>Phosphatase Activity (pNPP) A 200 μl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl₂ containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 50 pmol of Mth RNA Ligase incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.</p>	Pass
<p>Functional Testing (Adenylation) A 20 μl reaction in 1X 5' DNA Adenylation Reaction Buffer containing 100 pmol of 17 bp long 5'-ssDNA linker (5'-pCTGTAGGCACCATCAAT-NH₂-3'), 0.1 mM ATP, and 100 pmol of Mth RNA Ligase incubated for 1 hour at 65°C results in \geq95% 5'-ssDNA linker adenylated.</p>	Pass
<p>Exonuclease Activity (Radioactivity Release) A 50 μl reaction in 1X 5' DNA Adenylation Reaction Buffer containing 1 μg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 50 pmol of Mth RNA Ligase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p>	Pass

Assay Name/Specification	Lot # 10053873
<p>Endonuclease Activity (Nicking) A 50 µl reaction in 1X 5' DNA Adenylation Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 pmol of Mth RNA Ligase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p>RNase Activity Assay (4 Hour Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 50 pmol of Mth RNA Ligase is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p>* Individual Product Component Note Standard Quality Control Tests are performed for each component included in 5' DNA Adenylation Kit and meet the designated specifications.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.



Bo Wu
Production Scientist
11 Jul 2019



Michael Tonello
Packaging Quality Control Inspector
26 Nov 2019