


New England Biolabs Certificate of Analysis

Product Name: T7 Exonuclease
Catalog #: M0263S/L
Concentration: 10,000 units/ml
Unit Definition: One unit is defined as the amount of enzyme required to produce 1 nmol of acid-soluble deoxyribonucleotide in a total reaction volume of 50 µl in 30 minutes at 37°C in 1X NEBuffer 4 with 0.15 mM sonicated duplex [³H]-DNA.
Lot #: 0041712
Assay Date: 12/2017
Expiration Date: 12/2019
Storage Temp: -20°C
Storage Conditions: 10 mM Tris-HCl, 5 mM DTT, 0.1 mM EDTA, 50 % Glycerol, (pH 8.0 @ 25°C)
Specification Version: PS-M0263S/L v1.0
Effective Date: 23 May 2018

Assay Name/Specification (minimum release criteria)	Lot #0041712
Endonuclease Activity (Nicking) - A 50 µl reaction in NEBuffer 4 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 100 units of T7 Exonuclease incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) - T7 Exonuclease is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
RNase Activity (Extended Digestion) - A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 10 units of T7 Exonuclease is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
Single Stranded DNase Activity (FAM-Labeled Oligo) - A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 10 units of T7 Exonuclease incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.	Pass



Authorized by
Derek Robinson
23 May 2018



Inspected by
John Greci
03 Jan 2018

