

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: LongAmp® Taq DNA Polymerase

Catalog Number: M0323S
Concentration: 2,500 U/ml

Unit Definition: One unit is defined as the amount of enzyme that will incorporate 10

nmol of dNTP into acid insoluble material in 30 minutes at 75°C.

Packaging Lot Number: 10160439
Expiration Date: 07/2024
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 100 mM KCl, 1 mM DTT, 0.1 mM EDTA, 0.5 % Tween®

20 , 0.5 % IGEPAL® CA-630 , 50 % Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0323S/L v2.0

LongAmp® Taq DNA Polymerase Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0323SVIAL	LongAmp® Taq DNA Polymerase	10157641	Pass	
B0323SVIAL	LongAmp® Taq Reaction Buffer	10142614	Pass	

Assay Name/Specification	Lot # 10160439
qPCR DNA Contamination (E. coli Genomic) A minimum of 2.5 units of LongAmp® Taq DNA Polymerase is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass
PCR Amplification (30 kb Lambda DNA) A 25 µl reaction in LongAmp® Taq Reaction Buffer in the presence of 300 µM dNTPs and 0.4 µM primers containing 1 ng Lambda DNA with 2.5 units of LongAmp® Taq DNA Polymerase for 28 cycles of PCR amplification results in the expected 30 kb product.	Pass
PCR Amplification (30 kb Human Genomic DNA) A 25 μl reaction in LongAmp® Taq Reaction Buffer in the presence of 300 μM dNTPs and 0.4 μM primers containing 500 ng Human Genomic DNA with 2.5 units of LongAmp® Taq DNA Polymerase for 28 cycles of PCR amplification results in the expected 30 kb product.	Pass
RNase Activity (Extended Digestion)	Pass



M0323S / Lot: 10160439

Page 1 of 2

Assay Name/Specification	Lot # 10160439
A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of LongAmp® Taq DNA Polymerase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 2.5 units of LongAmp® Taq DNA Polymerase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Lea Antonpoulos

Production Scientist

27 Jul 2022 **4**

Michael Tonello

Packaging Quality Control Inspector

12 Oct 2022



M0323S / Lot: 10160439

Page 2 of 2