

## New England Biolabs Certificate of Analysis


**Product Name:** Exonuclease V (RecBCD)  
**Catalog Number:** M0345S  
**Concentration:** 10,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to produce 1 nmol of acid-soluble deoxyribonucleotide from double-stranded DNA in 30 minutes at 37°C in a total reaction volume of 50 µl.  
**Packaging Lot Number:** 10064165  
**Expiration Date:** 08/2021  
**Storage Temperature:** -20°C  
**Storage Conditions:** 50 mM Tris-HCl, 100 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 0.1% Triton®X-100, (pH 7.5 @ 25°C)  
**Specification Version:** PS-M0345S/L v1.0

Exonuclease V (RecBCD) Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
P0756SVIAL	Adenosine 5'-Triphosphate (ATP)	10058045	Pass
M0345SVIAL	Exonuclease V (RecBCD)	10051890	Pass
B7004SVIAL	NEBuffer™ 4	10061301	Pass

Assay Name/Specification	Lot # 10064165
<b>RNase Activity (Extended Digestion)</b> A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 10 units of Exonuclease V (RecBCD) is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
<b>Protein Purity Assay (SDS-PAGE)</b> Exonuclease V (RecBCD) is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in NEBuffer 4 supplemented with 1 mM ATP containing 1 µg of supercoiled PhiX174 RF I DNA and a minimum of 100 units of Exonuclease V (RecBCD) incubated for 4 hours at 37°C results in <10% loss in supercoiled DNA as determined by agarose gel electrophoresis.	Pass
<b>Endonuclease Activity (Nicked Double-Stranded DNA)</b>	Pass

Assay Name/Specification	Lot # 10064165
A 50 µl reaction in NEBuffer 4 supplemented with 1 mM ATP containing 1 µg of nicked PhiX174 RF II DNA and a minimum of 50 units of Exonuclease V (RecBCD) incubated for 4 hours at 37°C results in <10% loss in PhiX174 RF II DNA as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.



John Greci  
Production Scientist  
19 Aug 2019



Jay Minichiello  
Packaging Quality Control Inspector  
27 Jan 2020