NEW ENGLAND BioLabs Inc.
be INSPIRED drive DISCOVERY stay GENUINE
240 County Road
Ipswich, MA 01938-2723
Tel 978-927-5054
Fax 978-921-1350
www.neb.com
info@neb.com

New England Biolabs
Certificate of Analysis

Product Name:	*Antarctic Thermolabile UDG*
Catalog Number:	*M0372S*
Concentration:	*1,000 U/ml*
Unit Definition:	*One unit is defined as the amount of enzyme that catalyzes the release of 60 pmol of uracil per minute from double-stranded, uracil-containing DNA. Activity is measured by release of [3H] -uracil in a 50 µl reaction containing 0.2 µg DNA (104-105 cpm/µg) in 30 minutes at 37°C.*
Packaging Lot Number:	*10150213*
Expiration Date:	*02/2024*
Storage Temperature:	*-20°C*
Storage Conditions:	*50 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, (pH 7.4 @ 25°C)*
Specification Version:	*PS-M0372S/L v1.0*

Antarctic Thermolabile UDG Component List			
NEB Part Number	**Component Description**	**Lot Number**	**Individual QC Result**
M0372SVIAL	Antarctic Thermolabile UDG	10137676	**Pass**
B9014SVIAL	Standard Taq Reaction Buffer Pack	10143506	**Pass**

Assay Name/Specification	**Lot # 10150213**
Endonuclease Activity (Nicking) A 50 µl reaction in Standard Taq Reaction Buffer containing 1 µg of supercoiled PhiX174 RF I DNA and a minimum of 15 units of Antarctic Thermolabile UDG incubated for 4 hours at 37ºC results in <20% conversion to RFII as determined by agarose gel electrophoresis.	**Pass**
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in Standard Taq Reaction Buffer containing 1 µg of HindIII digested Lambda DNA and a minimum of 50 units of Antarctic Thermolabile UDG incubated for 16 hours at 37ºC results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	**Pass**
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of f-300 RNA transcript and a minimum of 1 unit of Antarctic Thermolabile UDG is incubated at 37ºC. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by	**Pass**

NEW ENGLAND BioLabs Inc.
be INSPIRED drive DISCOVERY stay GENUINE
240 County Road
Ipswich, MA 01938-2723
Tel 978-927-5054
Fax 978-921-1350
www.neb.com
info@neb.com

Assay Name/Specification	Lot # 10150213
gel electrophoresis using agarose gel electrophoresis.	
qPCR DNA Contamination (E. coli Genomic) A minimum of 1 unit of Antarctic Thermolabile UDG is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	**Pass**
Protein Purity Assay (SDS-PAGE) Antarctic Thermolabile UDG is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	**Pass**
Single Stranded DNase Activity (FAM-Labeled Oligo) A 50 µl reaction in NEBuffer 4 containing a 20 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 1 unit of Antarctic Thermolabile UDG incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.	**Pass**
DNase Activity (Labeled Oligo, 5' extension) A 50 µl reaction in NEBuffer 4 containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a 5' extension and a minimum of 1 unit of Antarctic Thermolabile UDG incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.	**Pass**
Double Stranded DNase Activity (Labeled Oligo) A 50 µl reaction in NEBuffer 4 containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a blunt end and a minimum of 1 unit of Antarctic Thermolabile UDG incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.	**Pass**
DNase Activity (Labeled Oligo, 3' extension) A 50 µl reaction in NEBuffer 4 containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a 3' extension and a minimum of 1 unit of Antarctic Thermolabile UDG incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.	**Pass**

This product has been tested and shown to be in compliance with all specifications.

NEW ENGLAND BioLabs Inc.
be INSPIRED drive DISCOVERY stay GENUINE
240 County Road
Ipswich, MA 01938-2723
Tel 978-927-5054
Fax 978-921-1350
www.neb.com
info@neb.com

Lauren Higgins
Production Scientist
25 Apr 2022

Erin Varney
Packaging Quality Control Inspector
25 Apr 2022