

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Quick CIP
Catalog Number: M0525S
Concentration: 5,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme that hydrolyzes 1 µmol

of p-Nitrophenyl Phosphate, PNPP in a total reaction volume of 1 ml

in 1 minute at 37°C.

Packaging Lot Number: 10067921 Expiration Date: 10/2021 Storage Temperature: -20°C

Storage Conditions: 25 mM Tris-HCl , 1 mM MgCl2 , 0.1 mM ZnCl2 , 50 % Glycerol, (pH 7.5

@ 25°C)

Specification Version: PS-M0525S/L v1.0

Quick CIP Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0525SVIAL	Quick CIP	10054876	Pass	
B7204SVIAL	CutSmart® Buffer	10068805	Pass	

Assay Name/Specification	Lot # 10067921
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart® Buffer containing 1 µg of supercoiled PhiX174 DNA and	Pass
a minimum of 50 units of Quick CIP incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA	Pass
and a minimum of 1 µl of Quick CIP is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis	
using gel electrophoresis using fluorescent detection.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 50 units of Quick CIP incubated	
for 4 hours at 37°C releases <0.1% of the total radioactivity.	
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in NEBuffer 4 containing 1 µg of PhiX174-HaeIII DNA and a minimum	



M0525S / Lot: 10067921

Page 1 of 2



Assay Name/Specification	Lot # 10067921
of 50 units of Quick CIP incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

Ana Egana

Production Scientist

24 Mar 2020

Mary Conlon

Packaging Quality Control Inspector

24 Mar 2020

