

New England Biolabs Certificate of Analysis

Product Name: *Histone H2A/H2B Dimer Human, Recombinant*
Catalog #: *M2508S*
Concentration: *20 µM*
Unit Definition: *N/A*
Lot #: *0031711*
Assay Date: *11/2017*
Expiration Date: *11/2018*
Storage Temp: *-20°C*
Storage Conditions: *2 M NaCl, 20 mM Tris-HCl, 1 mM DTT, 1 mM EDTA, (pH 8.0 @ 25°C)*
Specification Version: *PS-M2508S v1.0*
Effective Date: *25 Sep 2017*

Assay Name/Specification (minimum release criteria)	Lot #0031711
Endonuclease Activity (Nicking) - A 50 µl reaction in NEBuffer 2 containing 1 µg of supercoiled PhiX174 RF I DNA and a minimum of 10 µg of Histone H2A/H2B Dimer Human, Recombinant incubated for 4 hours at 37°C results in <10% conversion to RFII as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) - A 50 µl reaction in NEBuffer 2 containing 1 µg of a mixture of single and double-stranded [³ H] <i>E. coli</i> DNA and a minimum of 10 µg of Histone H2A/H2B Dimer Human, Recombinant incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Protease Activity (Histones) - A 12 µl reaction containing 7 µl of a standard mixture of proteins and a minimum of 10 µg of Histone H2A/H2B Dimer Human, Recombinant incubated for 4 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.	Pass
Protein Purity Assay (SDS-PAGE) - Histone H2A/H2B Dimer Human, Recombinant is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass



Authorized by
Derek Robinson
25 Sep 2017



Inspected by
Fana Mersha
20 Nov 2017

