240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Lambda DNA (N6-methyladenine-free)

 Catalog #:
 N3013S/L

 Concentration:
 500 μg/ml

 Lot #:
 0491603

 Assay Date:
 03/2016

 Expiration Date:
 03/2018

 Storage Temp:
 -20°C

Storage Conditions: 10 mM Tris-HCl (pH 8.0), 1 mM EDTA

Specification Version: PS-N3013S/L v1.0
Effective Date: 20 Nov 2014

Assay Name/Specification (minimum release criteria)	Lot #0491603
A260/A280 Assay - The ratio of UV absorption of Lambda DNA (N6-methyladenine-free) at 260 and 280 nm is between 1.8 and 2.0.	Pass
DNA Concentration (A260) - The concentration of Lambda DNA (N6-methyladenine-free) is between 500 and 550 μg/ml as determined by UV absorption at 260 nm.	Pass
Electrophoretic Pattern (Linear DNA) - The banding pattern of Lambda DNA (N6-methyladenine-free) on a 1.2% agarose gel is evaluated against a control lot for sharpness and relative intensity as determined by gel electrophoresis using Ethidium Bromide.	Pass
Non-Specific DNase Activity (DNA, 16 hour) - A 50 μl reaction in 1X NEBuffer 2 containing 2.5 μg of Lambda DNA (N6-methyladenine-free) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Restriction Digest (Correct Pattern) - A 50 μl reaction in NEBuffer 2.1 containing 2.5 μg of Lambda DNA (N6-methyladenine-free) DNA and 20 units of HindIII incubated for 1 hour at 37°C produces the expected pattern of DNA fragments as determined by agarose gel electrophoresis.	Pass
Restriction Digest (Dam Resistant) - A 50 μl reaction in CutSmart TM Buffer containing 2.5 μg of Lambda DNA (N6-methyladenine-free) and a minimum of 20 units of DpnI incubated for 1 hour at 37°C results in no detectable digestion of the DNA as determined by agarose gel electrophoresis.	Pass
Restriction Digest (Dam Sensitive) - A 50 µl reaction in NEBuffer DpnII containing 2.5 µg of Lambda DNA (N6-methyladenine-free) DNA and a minimum of 10 units of DpnII incubated for 1 hour at 37°C results in complete digestion of the DNA as determined by agarose gel electrophoresis.	Pass

Authorized by Derek Robinson 20 Nov 2014







Inspected by

Vanessa Mathieu-Sheltry

23 Mar 2016