

New England Biolabs Certificate of Analysis

Product Name: Endo H
Catalog Number: P0702S
Concentration: 500,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to remove > 95% of the carbohydrate from 10 µg of denatured RNase B in 1 hour at 37°C in a total reaction volume of 10 µl (10 NEB units = 1 IUB milliunit).
Packaging Lot Number: 10096798
Expiration Date: 02/2023
Storage Temperature: -20°C
Storage Conditions: 50 mM NaCl, 20 mM Tris-HCl, 5 mM EDTA, (pH 7.5 @ 25°C)
Specification Version: PS-P0702S/L v1.0

Endo H Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
P0702SVIAL	Endo H	10096799	Pass
B1720SVIAL	10X Glycobuffer 3	10082734	Pass
B1704SVIAL	Glycoprotein Denaturing Buffer	10090539	Pass

Assay Name/Specification	Lot # 10096798
Protein Purity Assay (SDS-PAGE) Endo H is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Protease Activity (SDS-PAGE) A 20 µl reaction in 1X Glyco Buffer 3 containing 24 µg of a standard mixture of proteins and a minimum of 5,000 units of Endo H incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.	Pass
Glycosidase Activity (β1-4 Galactosidase) A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-4GlcNAcβ1-3Galβ1-4Glc -AMC) and 5,000 units of Endo H incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β1-3 Galactosidase)	Pass

Assay Name/Specification	Lot # 10096798
<p>A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC) and 5,000 units of Endo H incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	
<p>Glycosidase Activity (β-N-Acetylgalactosaminidase) A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled β-N-Acetylgalactosaminidase substrate (GalNAcβ1-4Galβ1-4Glc-AMC) and 5,000 units of Endo H incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-Mannosidase) A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled β-Mannosidase substrate (Manβ1-4Manβ1-4Man-AMC) and 5,000 units of Endo H incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-Glucosidase) A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α-Glucosidase substrate (Glcα1-6Glcα1-4Glc-AMC) and 5,000 units of Endo H incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-6 Mannosidase) A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α-Mannosidase substrate (Manα1-6Manα1-6(Manα1-3)Man-AMC) and 5,000 units of Endo H incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-Neuraminidase) A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α-Neuraminidase substrate (Neu5Acα2-3Galβ1-3GlcNAcβ1-3Galβ1-4Glc-AMC) and 5,000 units of Endo H incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-N-Acetylgalactosaminidase) A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α-N-Acetylgalactosaminidase substrate (GalNAcα1-3(Fucα1-2)Galβ1-4Glc-AMC) and 5,000 units of Endo H incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-Xylosidase) A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled</p>	Pass

Assay Name/Specification	Lot # 10096798
<p>β-Xylosidase substrate (Xylβ1-4Xylβ1-4Xylβ1-4Xyl-AMC) and 5,000 units of Endo H incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	
<p>Glycosidase Activity (β-N-Acetylglucosaminidase) A 10 μl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled β-N-Acetylglucosaminidase substrate (GlcNAcβ1-4GlcNAcβ1-4GlcNAc-AMC) and 5,000 units of Endo H incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-3 Galactosidase) A 10 μl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α-Galactosidase substrate (Galα1-3Galβ1-4GlcNAc-AMC) and 5,000 units of Endo H incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-3 Fucosidase) A 10 μl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α-Fucosidase substrate (Fucα1-3Galβ1-4GlcNAcβ1-3Galβ1-4Glc-AMC) and 5,000 units of Endo H incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-6 Galactosidase) A 10 μl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α-Galactosidase substrate (Galα1-6Galα1-6Glcα1-2Fru-AMC) and 5,000 units of Endo H incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-3 Mannosidase) A 10 μl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α-Mannosidase substrate (Manα1-3Manβ1-4GlcNAc-AMC) and 5,000 units of Endo H incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-2 Fucosidase) A 10 μl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled α-Fucosidase substrate (Fucα1-2Galβ1-4Glc-AMC) and 5,000 units of Endo H incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (PNGase F) A 10 μl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled PNGase F substrate (Fluoresceinated fetuin triantennary) and 5,000 units of Endo H incubated</p>	Pass

Assay Name/Specification	Lot # 10096798
<p>for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> <p>Glycosidase Activity (Endo F2, F3) A 10 µl reaction in Glyco Buffer 3 containing 1 nM of fluorescently-labeled Endo F2, F3 substrate (Dansylated fibrinogen biantennary) and 5,000 units of Endo H incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<p>Pass</p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Alicia Bielik
Production Scientist
12 Feb 2021



Josh Hersey
Packaging Quality Control Inspector
12 Feb 2021