

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Bacteroides Heparinase I

Catalog Number: P0735S
Concentration: 12,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme that will liberate 1.0

µmol unsaturated oligosaccharides from porcine mucosal heparin per

minute at 30°C and pH 7.0 in a total reaction volume of 100 μl.

Packaging Lot Number: 10226178
Expiration Date: 12/2024
Storage Temperature: -80°C

Storage Conditions: 100 mM NaCl, 20 mM Tris-HCl, 1 mM EDTA, 5 mM CaCl2, (pH 7.5 @ 25°C)

Specification Version: PS-P0735S/L v1.0

Bacteroides Heparinase I Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
P0735SVIAL	Bacteroides Heparinase I	10221747	Pass	
B0735SVIAL	Bacteroides Heparinase Reaction Buffer (10X)	10204911	Pass	

Assay Name/Specification	Lot # 10226178
Glycosidase Activity (β-N-Acetylgalactosaminidase) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled β-N-Acetylgalactosaminidase substrate (GalNAc $\beta$ 1-4Gal $\beta$ 1-4Glc-AMC) and 24 units of Bacteroides Heparinase I incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β-N-Acetylglucosaminidase) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled β-N-Acetylglucosaminidase substrate (GlcNAcβ1-4GlcNAcβ1-4GlcNAc-AMC) and 24 units of Bacteroides Heparinase I incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β1-3 Galactosidase) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC) and 24 units of Bacteroides Heparinase I incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.	Pass



P0735S / Lot: 10226178

Page 1 of 3

Assay Name/Specification	Lot # 10226178
Glycosidase Activity (β1-4 Galactosidase) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-4GlcNAcβ1-3Galβ1-4Glc -AMC) and 24 units of Bacteroides Heparinase I incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.	Pass
Protease Activity (SDS-PAGE) A 20 µl reaction in 1X Heparinase Reaction Buffer containing 24 µg of a standard mixture of proteins and a minimum of 120 units of Bacteroides Heparinase I incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.	Pass
Protein Purity Assay (SDS-PAGE) Bacteroides Heparinase I is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Sulfatase Activity (2-O) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled 2-O-Sulfatase substrate (ΔUA2S-(1-4)-GlcNS6S-AMC) and 24 units of Bacteroides Heparinase I incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.	Pass
Sulfatase and Uronidase Activity (N,6-O) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled N,6-O-Sulfatase substrate (ΔUA-(1-4)-GlcNS6S-AMC) and 24 units of Bacteroides Heparinase I incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



P0735S / Lot: 10226178

Page 2 of 3

Production Scientist 20 Dec 2023

Josh Hersey Packaging Quality Control Inspector

24 Jan 2024

P0735S / Lot: 10226178

Page 3 of 3