

New England Biolabs Certificate of Analysis

Product Name: *Bacteroides Heparinase II*
Catalog #: *P0736S/L*
Concentration: *4,000 units/ml*
Unit Definition: *One unit is defined as the amount of enzyme that will liberate 1.0 μmol unsaturated oligosaccharides from porcine mucosal heparin per minute at 30°C and pH 7.0 in a total reaction volume of 100 μl.*
Lot #: *0031803*
Assay Date: *03/2018*
Expiration Date: *3/2019*
Storage Temp: *-80°C*
Storage Conditions: *100 mM NaCl, 20 mM Tris-HCl, 1 mM EDTA, 5 mM CaCl₂, (pH 7.5 @ 25°C)*
Specification Version: *PS-P0736S/L v1.0*
Effective Date: *16 Feb 2016*


Assay Name/Specification (minimum release criteria)	Lot #0031803
Glycosidase Activity (β1-3 Galactosidase) - A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC) and 8 units of <i>Bacteroides</i> Heparinase II incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β1-4 Galactosidase) - A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-4GlcNAcβ1-3Galβ1-4Glc-AMC) and 8 units of <i>Bacteroides</i> Heparinase II incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β-N-Acetylgalactosaminidase) - A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled β-N-Acetylgalactosaminidase substrate (GalNAcβ1-4Galβ1-4Glc-AMC) and 8 units of <i>Bacteroides</i> Heparinase II incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β-N-Acetylglucosaminidase) - A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled β-N-Acetylglucosaminidase substrate (GlcNAcβ1-4GlcNAcβ1-4GlcNAc-AMC) and 8 units of <i>Bacteroides</i> Heparinase II incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.	Pass

New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0031803
Protease Activity (SDS-PAGE) - A 20 µl reaction in 1X Heparinase Reaction Buffer containing 24 µg of a standard mixture of proteins and a minimum of 20 units of <i>Bacteroides</i> Heparinase II incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.	Pass
Protein Purity Assay (SDS-PAGE) - <i>Bacteroides</i> Heparinase II is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Sulfatase Activity (2γ) - A 10 µl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled 2-O-Sulfatase substrate (ΔUA2S-(1-4)-GlcNS6S-AMC) and 8 units of <i>Bacteroides</i> Heparinase II incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Sulfatase and Uronidase Activity (N,6γ) - A 10 µl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled N,6-O-Sulfatase substrate (ΔUA-(1-4)-GlcNS6S-AMC) and 8 units of <i>Bacteroides</i> Heparinase II incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.	Pass



Authorized by
Derek Robinson
16 Feb 2016



Inspected by
Brad Landgraf
16 Mar 2018

