

## New England Biolabs Certificate of Analysis

**Product Name:** SpeI  
**Catalog #:** R0133S/L  
**Concentration:** 10,000 units/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of pXba-XbaI DNA in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Lot #:** 0321511  
**Assay Date:** 11/2015  
**Expiration Date:** 11/2017  
**Storage Temp:** -20°C  
**Storage Conditions:** 250 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 0.15% Triton X-100, 200 µg/ml BSA  
**Specification Version:** PS-R0133S/L v1.0  
**Effective Date:** 07 Jun 2013

Assay Name/Specification (minimum release criteria)	Lot #0321511
<b>Blue-White Screening (Terminal Integrity)</b> - A sample of LITMUS28 vector linearized with a 10-fold excess of SpeI, religated and transformed into an <i>E. coli</i> strain expressing the LacZ beta fragment gene results in <1% white colonies.	<b>Pass</b>
<b>Endonuclease Activity (Nicking)</b> - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 Units of SpeI incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	<b>Pass</b>
<b>Exonuclease Activity (Radioactivity Release)</b> - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 50 units of SpeI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	<b>Pass</b>
<b>Ligation and Recutting (Terminal Integrity)</b> - After a 20-fold over-digestion of Adenovirus-2 DNA with SpeI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with SpeI.	<b>Pass</b>
<b>Non-Specific DNase Activity (16 Hour)</b> - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pXba-XbaI digested DNA and a minimum of 50 units of SpeI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	<b>Pass</b>
<b>Protein Purity Assay (SDS-PAGE)</b> - SpeI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	<b>Pass</b>

---


## New England Biolabs Certificate of Analysis

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.



---

Authorized by  
Derek Robinson  
07 Jun 2013



---

Inspected by  
Terry Petronzio  
02 Nov 2015

