

## New England Biolabs Certificate of Analysis

*Product Name:* TaqαI  
*Catalog #:* R0149S/L  
*Concentration:* 20,000 units/ml  
*Unit Definition:* One unit is defined as the amount of enzyme required to digest 1 αg of Lambda DNA in 1 hour at 65°C in a total reaction volume of 50 αl.  
*Lot #:* 0541405  
*Assay Date:* 05/2014  
*Expiration Date:* 5/2016  
*Storage Temp:* -20 °C  
*Storage Conditions:* 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 μg/ml BSA  
*Specification Version:* PS-R0149S/L v1.0  
*Effective Date:* 24 Apr 2014

Assay Name/Specification (minimum release criteria)	Lot #0541405
<b>Exonuclease Activity (Radioactivity Release)</b> - A 50 μl reaction in CutSmart™ Buffer containing 1 μg of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 200 units of TaqαI incubated for 4 hours at 65°C releases <0.1% of the total radioactivity.	<b>Pass</b>
<b>Ligation and Recutting (Terminal Integrity)</b> - After a 20-fold over-digestion of Lambda DNA DNA with TaqαI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 25°C. Of these ligated fragments, >95% can be recut with TaqαI.	<b>Pass</b>
<b>Non-Specific DNase Activity (16 Hour)</b> - A 50 μl reaction in CutSmart™ Buffer containing 1 μg of Lambda DNA DNA and a minimum of 200 Units of TaqαI incubated for 16 hours at 65°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	<b>Pass</b>

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.



Authorized by  
Derek Robinson  
24 Apr 2014



Inspected by  
David Hough  
24 Apr 2014

