

## New England Biolabs Certificate of Analysis

**Product Name:** TaqI-v2  
**Catalog Number:** R0149S  
**Concentration:** 20,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 65°C in a total reaction volume of 50 µl.  
**Packaging Lot Number:** 10079895  
**Expiration Date:** 03/2022  
**Storage Temperature:** -20°C  
**Storage Conditions:** 300 mM NaCl , 10 mM Tris-HCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol , 500 µg/ml BSA, (pH 7.4 @ 25°C)  
**Specification Version:** PS-R0149S/L v2.0

TaqI-v2 Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0149SVIAL	TaqI-v2	10068121	Pass
B7204SVIAL	CutSmart® Buffer	10079734	Pass

Assay Name/Specification	Lot # 10079895
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 200 units of TaqI-v2 incubated for 4 hours at 65°C releases <0.1% of the total radioactivity.	Pass
<b>Functional Testing (15 minute Digest)</b> A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and 1 µl of TaqI-v2 incubated for 15 minutes at 65°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 20-fold over-digestion of Lambda DNA with TaqI-v2, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 25°C. Of these ligated fragments, >95% can be recut with TaqI-v2.	Pass
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and a minimum of 200 units of TaqI-v2 incubated for 16 hours at 65°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass


Assay Name/Specification	Lot # 10079895
<p><b>Protein Purity Assay (SDS-PAGE)</b> TaqI-v2 is <math>\geq</math> 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	<b>Pass</b>
<p><b>RNase Activity (Extended Digestion)</b> A 10 <math>\mu</math>l reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 <math>\mu</math>l of TaqI-v2 is incubated at 37°C. After incubation for 16 hours, &gt;90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Penghua Zhang  
Production Scientist  
10 Sep 2020



Michael Tonello  
Packaging Quality Control Inspector  
10 Sep 2020