

New England Biolabs Certificate of Analysis

Product Name: TaqαI
Catalog #: R0149T/M
Concentration: 100,000 units/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 αg of Lambda DNA in 1 hour at 65°C in a total reaction volume of 50 αl.
Lot #: 0541604
Assay Date: 04/2016
Expiration Date: 04/2018
Storage Temp: -20°C
Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 μg/ml BSA
Specification Version: PS-R0149T/M v1.0
Effective Date: 28 Jun 2013

Assay Name/Specification (minimum release criteria)	Lot #0541604
Exonuclease Activity (Radioactivity Release) - A 50 μl reaction in CutSmart™ Buffer containing 1 μg of a mixture of single and double-stranded [³ H] <i>E. coli</i> DNA and a minimum of 200 units of TaqαI incubated for 4 hours at 65°C releases <0.1% of the total radioactivity.	Pass
Ligation and Recutting (Terminal Integrity) - After a 20-fold over-digestion of Lambda DNA DNA with TaqαI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 25°C. Of these ligated fragments, >95% can be recut with TaqαI.	Pass
Non-Specific DNase Activity (16 Hour) - A 50 μl reaction in CutSmart™ Buffer containing 1 μg of Lambda DNA DNA and a minimum of 200 Units of TaqαI incubated for 16 hours at 65°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.



Authorized by
Derek Robinson
28 Jun 2013



Inspected by
Jianying Luo
15 Apr 2016

