

New England Biolabs Certificate of Analysis

Product Name: NarI
Catalog Number: R0191S
Concentration: 5,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg pXba DNA in 1 hour at 37°C in a total reaction volume of 50 µl.
Lot Number: 10050285
Expiration Date: 07/2021
Storage Temperature: -20°C
Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA
Specification Version: PS-R0191S/L v2.0

NarI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0191SVIAL	NarI	10050287	Pass
B7204SVIAL	CutSmart® Buffer	10046087	Pass

Assay Name/Specification	Lot # 10050285
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart Buffer containing 1 µg of supercoiled LITMUS28i DNA and a minimum of 50 units of NarI incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 50 units of NarI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of pXba DNA with NarI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with NarI.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart Buffer containing 1 µg of pXba DNA and a minimum of 50 units of NarI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.



Stephanie Cornelio
Production Scientist
23 Jul 2019



Jay Minichiello
Packaging Quality Control Inspector
14 Aug 2019