NEW ENGLAND BioLabs Inc.
be INSPIRED drive DISCOVERY stay GENUINE
240 County Road
Ipswich, MA 01938-2723
Tel 978-927-5054
Fax 978-921-1350
www.neb.com
info@neb.com

New England Biolabs
Certificate of Analysis

Product Name:	*BstYI*
Catalog Number:	*R0523S*
Concentration:	*10,000 U/ml*
Unit Definition:	*One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 60°C in a total reaction volume of 50 µl.*
Packaging Lot Number:	*10132334*
Expiration Date:	*12/2023*
Storage Temperature:	*-20°C*
Storage Conditions:	*50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA*
Specification Version:	*PS-R0523S/L v1.0*

BstYI Component List			
NEB Part Number	**Component Description**	**Lot Number**	**Individual QC Result**
R0523SVIAL	BstYI	10132333	**Pass**
B6002SVIAL	NEBuffer™ r2.1	10103928	**Pass**

Assay Name/Specification	**Lot # 10132334**
Protein Purity Assay (SDS-PAGE) BstYI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	**Pass**
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 2.1 containing 1 µg of Lambda DNA and a minimum of 30 Units of BstYI incubated for 16 hours at 60ºC results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	**Pass**
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of Lambda DNA with BstYI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16ºC. Of these ligated fragments, >95% can be recut with BstYI.	**Pass**
Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 2.1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 10 Units of BstYI incubated for 4 hours at 60ºC results in <50% conversion to the nicked form as determined by agarose gel electrophoresis.	**Pass**

NEW ENGLAND BioLabs Inc.
be INSPIRED drive DISCOVERY stay GENUINE
240 County Road
Ipswich, MA 01938-2723
Tel 978-927-5054
Fax 978-921-1350
www.neb.com
info@neb.com

Assay Name/Specification	Lot # 10132334
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 2.1 containing 1 µg of a mixture of single and double-stranded [^{3}H] E. coli DNA and a minimum of 50 units of BstYI incubated for 4 hours at 60ºC releases <0.1% of the total radioactivity.	**Pass**

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang
Production Scientist
04 Jan 2022

Michael Tonello
Packaging Quality Control Inspector
04 Jan 2022