

## New England Biolabs Certificate of Analysis

**Product Name:** SgrAI  
**Catalog Number:** R0603S  
**Concentration:** 10,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of LambdaDNA in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Packaging Lot Number:** 10069181  
**Expiration Date:** 03/2022  
**Storage Temperature:** -20°C  
**Storage Conditions:** 100 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA  
**Specification Version:** PS-R0603S/L v1.0

SgrAI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0603SVIAL	SgrAI	10069180	Pass
B7204SVIAL	CutSmart® Buffer	10071078	Pass

Assay Name/Specification	Lot # 10069181
<p><b>Non-Specific DNase Activity (16 hour)</b>            A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 10 Units of SgrAI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. NOTE: although no nuclease degradation is detected under these conditions, extended incubations and/or high concentrations of this enzyme may result in star activity. See the product FAQ for recommended reaction conditions for this enzyme.</p>	Pass
<p><b>Ligation and Recutting (Terminal Integrity)</b>            After a 2-fold over-digestion of Lambda DNA with SgrAI, &gt;95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, &gt;95% can be recut with SgrAI.</p>	Pass
<p><b>Exonuclease Activity (Radioactivity Release)</b>            A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] E. coli DNA and a minimum of 50 units of SgrAI incubated for 4 hours at 37°C releases &lt;0.1% of the total radioactivity.</p>	Pass
<p><b>Endonuclease Activity (Nicking)</b></p>	Pass

Assay Name/Specification	Lot # 10069181
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 30 Units of SgrAI incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

  
\_\_\_\_\_  
Penghua Zhang  
Production Scientist  
22 Apr 2020

  
\_\_\_\_\_  
Jay Minichiello  
Packaging Quality Control Inspector  
22 Apr 2020