NEW ENGLAND BioLabs Inc.
be INSPIRED drive DISCOVERY stay GENUINE
240 County Road
Ipswich, MA 01938-2723
Tel 978-927-5054
Fax 978-921-1350
www.neb.com
info@neb.com

# New England Biolabs
# Certificate of Analysis

| | |
|---|---|
| *Product Name:* | *SfoI* |
| *Catalog Number:* | *R0606L* |
| *Concentration:* | *10,000 U/ml* |
| *Unit Definition:* | *One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA (HindIII digest) in rCutSmart Buffer in 1 hour at 37°C in a total reaction volume of 50 µl.* |
| *Packaging Lot Number:* | *10167915* |
| *Expiration Date:* | *10/2024* |
| *Storage Temperature:* | *-20°C* |
| *Storage Conditions:* | *10 mM Tris-HCl, 200 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml rAlbumin (pH 7.4 @ 25°C)* |
| *Specification Version:* | *PS-R0606S/L v2.0* |

| **SfoI Component List** | | | |
|---|---|---|---|
| **NEB Part Number** | **Component Description** | **Lot Number** | **Individual QC Result** |
| R0606LVIAL | SfoI | 10167914 | **Pass** |
| B6004SVIAL | rCutSmart™ Buffer | 10163560 | **Pass** |

| **Assay Name/Specification** | **Lot # 10167915** |
|---|---|
| **Endonuclease Activity (Nicking)**<br>A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of supercoiled LITMUS28i DNA and a minimum of 30 units of SfoI incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis. | **Pass** |
| **Exonuclease Activity (Radioactivity Release)**<br>A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ $^3$H] E. coli DNA and a minimum of 100 units of SfoI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | **Pass** |
| **Non-Specific DNase Activity (16 Hour)**<br>A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda-HindIII DNA and a minimum of 30 units of SfoI incubated for 16 hours at 37ºC results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | **Pass** |
| **Ligation and Recutting (Terminal Integrity)**<br>After a 10-fold over-digestion of Lambda-HindIII DNA with SfoI, >95% of the DNA | **Pass** |

NEW ENGLAND BioLabs Inc.
be INSPIRED drive DISCOVERY stay GENUINE
240 County Road
Ipswich, MA 01938-2723
Tel 978-927-5054
Fax 978-921-1350
www.neb.com
info@neb.com

| Assay Name/Specification | Lot # 10167915 |
|---|---|
| fragments can be ligated with T4 DNA ligase in 16 hours at 16ºC. Of these ligated fragments, >95% can be recut with SfoI. | |
| **Protein Purity Assay (SDS-PAGE)**<br>SfoI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection. | **Pass** |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

YunJie Sun
Production Scientist
25 Oct 2022

Michael Tonello
Packaging Quality Control Inspector
27 Oct 2022