

New England Biolabs Certificate of Analysis

Product Name: Nb.BbvCI
Catalog #: R0631S/L
Concentration: 10,000 units/ml
Unit Definition: One unit is defined as the amount of enzyme required to convert 1 µg of supercoiled pUB DNA to open circular form in 1 hour at 37°C in a total reaction volume of 50 µl.
Lot #: 0031610
Assay Date: 10/2016
Expiration Date: 10/2018
Storage Temp: -20°C
Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA
Specification Version: PS-R0631S/L v2.0
Effective Date: 31 Jan 2014

Assay Name/Specification (minimum release criteria)	Lot #0031610
Exonuclease Activity (Radioactivity Release) - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³ H] <i>E. coli</i> DNA and a minimum of 30 units of Nb.BbvCI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Non-Specific DNase Activity (16 hour) - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pUB DNA and a minimum of 10 units of Nb.BbvCI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. NOTE: although no nuclease degradation is detected under these conditions, extended incubations and/or high concentrations of this enzyme may result in star activity. See the product FAQ for recommended reaction conditions for this enzyme.	Pass

* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.



Authorized by
Derek Robinson
31 Jan 2014



Inspected by
Jianying Luo
25 Oct 2016

