

New England Biolabs Certificate of Analysis


Product Name: CviKI-1
Catalog #: R0710S/L
Concentration: 5,000 units/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of pBR322 DNA in 1 hour at 37°C in a total reaction volume of 50 µl.
Lot #: 0031704
Assay Date: 04/2017
Expiration Date: 4/2019
Storage Temp: -20°C
Storage Conditions: 50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA
Specification Version: PS-R0710S/L v1.0
Effective Date: 30 Sep 2013

Assay Name/Specification (minimum release criteria)	Lot #0031704
<p>Ligation and Recutting (Terminal Integrity) - After a 5-fold over-digestion of pBR322 DNA with CviKI-1, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with CviKI-1.</p>	Pass
<p>Non-Specific DNase Activity (16 hour) - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pBR322 DNA and a minimum of 5 Units of CviKI-1 incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. NOTE: although no nuclease degradation is detected under these conditions, extended incubations and/or high concentrations of this enzyme may result in star activity. See the product FAQ for recommended reaction conditions for this enzyme.</p>	Pass

* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.



Authorized by
Derek Robinson
30 Sep 2013



Inspected by
Penghua Zhang
11 Apr 2017

