

## New England Biolabs Certificate of Analysis

**Product Name:** BsmBI-v2  
**Catalog Number:** R0739L  
**Concentration:** 10,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 55°C in a total reaction volume of 50 µl.  
**Packaging Lot Number:** 10226010  
**Expiration Date:** 09/2025  
**Storage Temperature:** -20°C  
**Storage Conditions:** 300 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 500 µg/ml BSA, (pH 7.4 @ 25°C)  
**Specification Version:** PS-R0739S/L v1.0


BsmBI-v2 Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0739LVIAL	BsmBI-v2	10206380	Pass
B7024AVIAL	Gel Loading Dye, Purple (6X)	10221467	Pass
B6003SVIAL	NEBuffer™ r3.1	10221488	Pass

Assay Name/Specification	Lot # 10226010
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in NEBuffer 3.1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 10 units of BsmBI-v2 incubated for 4 hours at 55°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 50 units of BsmBI-v2 incubated for 4 hours at 55°C releases <0.1% of the total radioactivity.	Pass
<b>Functional Testing (15 minute Digest)</b> A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and 1 µl of BsmBI-v2 incubated for 15 minutes at 55°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 10-fold over-digestion of Lambda DNA with BsmBI-v2, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated	Pass

Assay Name/Specification	Lot # 10226010
fragments, >95% can be recut with BsmBI-v2.	
<p><b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 10 units of BsmBI-v2 incubated for 16 hours at 55°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	<b>Pass</b>
<p><b>Protein Purity Assay (SDS-PAGE)</b> BsmBI-v2 is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



YunJie Sun  
Production Scientist  
13 Sep 2023



Michael Tonello  
Packaging Quality Control Inspector  
25 Jan 2024