

New England Biolabs Certificate of Analysis

Product Name: NsiI-HF[®]
Catalog #: R3127S/L
Concentration: 20,000 units/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.
Lot #: 0041612
Assay Date: 12/2016
Expiration Date: 12/2018
Storage Temp: -20°C
Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 500 µg/ml BSA
Specification Version: PS-R3127S/L v1.0
Effective Date: 08 Aug 2016

| Assay Name/Specification (minimum release criteria) | Lot #0041612 |
|---|--------------|
| Endonuclease Activity (Nicking) - A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of supercoiled ΦX174 DNA and a minimum of 100 units of NsiI-HF incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Exonuclease Activity (Radioactivity Release) - A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of a mixture of single and double-stranded [³ H] <i>E. coli</i> DNA and a minimum of 100 units of NsiI-HF incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass |
| Functional Test (15 minute Digest) - A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of Lambda DNA and 1 µl of NsiI-HF incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis. | Pass |
| Ligation and Recutting (Terminal Integrity) - After a 20-fold over-digestion of Lambda DNA with NsiI-HF, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with NsiI-HF. | Pass |
| Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of Lambda DNA and a minimum of 100 units of NsiI-HF incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |
| Protein Purity Assay (SDS-PAGE) - NsiI-HF is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection. | Pass |



New England Biolabs Certificate of Analysis

* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.



Authorized by
Derek Robinson
08 Aug 2016





Inspected by
Anthony Francis
27 Dec 2016