

## New England Biolabs Product Specification

<b>Product Name:</b>	<i>Bst</i> DNA Polymerase, Full Length
<b>Catalog #:</b>	M0328S/V
<b>Concentration:</b>	5,000 units/ml
<b>Unit Definition:</b>	One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTP into acid insoluble material in 30 minutes at 65°C.
<b>Shelf Life:</b>	24 months
<b>Storage Temp:</b>	-20°C
<b>Storage Conditions:</b>	10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 0.1 % Triton®X-100, 50 % Glycerol, (pH 7.1 @ 25°C)
<b>Specification Version:</b>	PS-M0328S/V v3.0
<b>Effective Date:</b>	17 Mar 2022

### Assay Name/Specification (minimum release criteria)

**Endonuclease Activity (Nicking)** - A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 units of *Bst* DNA Polymerase, Full Length incubated for 4 hours at 37°C and 65°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

**Non-Specific DNase Activity (16 Hour)** - A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 5 units of *Bst* DNA Polymerase, Full Length incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

**Phosphatase Activity (pNPP)** - A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl<sub>2</sub> containing 2.5 mM *p*-Nitrophenyl Phosphate (pNPP) and a minimum of 100 units *Bst* DNA Polymerase, Full Length incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.

**Protein Purity Assay (SDS-PAGE)** - *Bst* DNA Polymerase, Full Length is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.

**qPCR DNA Contamination (*E. coli* Genomic)** - A minimum of 5 units of *Bst* DNA Polymerase, Full Length is screened for the presence of *E. coli* genomic DNA using SYBR® Green qPCR with primers specific for the *E. coli* 16S rRNA locus. Results are quantified using a standard curve generated from purified *E. coli* genomic DNA. The measured level of *E. coli* genomic DNA contamination is ≤ 1 *E. coli* genome.



---

## New England Biolabs Product Specification

Assay Name/Specification (minimum release criteria)
---

RNase Activity Assay - A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 5 units of <i>Bst</i> DNA Polymerase, Full Length is incubated at 37°C. After incubation for 4 hours, the substrate RNA is assessed by gel electrophoresis using fluorescent detection and compared to the product's RNase QC Standard resulting in no additional non-specific nuclease degradation.
--

One or more products referenced in this document may be covered by a 3rd-party trademark.  
Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Date 17 Mar 2022

---

Derek Robinson  
Director, Quality Control

