

## New England Biolabs Product Specification

*Product Name:* Internal Control LAMP Primer Mix (rActin)  
*Catalog #:* S0164B  
*Concentration:* 10X Concentrate  
*Shelf Life:* 24 months  
*Storage Temp:* -20°C  
*Composition (1X):* Proprietary  
*Specification Version:* PS-S0164B v1.0  
*Effective Date:* 15 Jul 2020

### Assay Name/Specification (minimum release criteria)

**Endonuclease Activity (Nicking)** - A 50 µl reaction in NEBuffer 2 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 5 µl of Internal Control LAMP Primer Mix (rActin) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

**Non-Specific DNase Activity (16 Hour)** - A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 5 µl of Internal Control LAMP Primer Mix (rActin) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

**Phosphatase Activity (pNPP)** - A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl<sub>2</sub> containing 2.5 mM *p*-Nitrophenyl Phosphate (pNPP) and a minimum of 20 µl of Internal Control LAMP Primer Mix (rActin) incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.

**RNase Activity (Extended Digestion)** - A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Internal Control LAMP Primer Mix (rActin) is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.

*"One or more products referenced in this document may be covered by a 3rd-party trademark.  
Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information."*



Date 15 Jul 2020

Derek Robinson  
Director of Quality Control

